

Entropy bounds and field equations

Alessandro Pesci*

INFN-Bologna, Via Irnerio 46, I-40126 Bologna, Italy

For general metric theories of gravity, we compare the approach that describes-derives the field equations of gravity as a thermodynamic identity with the one which looks at them from entropy bounds. The comparison is made through the consideration of the matter entropy flux across (Rindler) horizons, studied by making use of the notion of a limiting thermodynamic scale l^* of matter, previously introduced in the context of entropy bounds.

In doing this: i) a bound for the entropy of any lump of matter with a given energy-momentum tensor T_{ab} is considered, in terms of a quantity which is independent of the theory of gravity we use; this quantity is the variation of the Clausius entropy of a suitable horizon when the element of matter crosses it; ii) by making use of the equations of motion of the theory, the same quantity is then expressed as the variation of Wald's entropy of that horizon (and this leads to a generalized form of the generalized covariant entropy bound, applicable to general diffeomorphism-invariant theories of gravity); iii) a notion of l^* for horizons, and an expression for it, is given.

PACS numbers: 04.20.Cv, 04.50.-h, 04.60.-m, 04.70.Dy

I. MOTIVATION

The thermodynamic description of the field equations of gravity was first performed by Jacobson in [1], where these equations were shown to be equivalent to a thermodynamic identity. In [2, 3], Padmanabhan has shown that this identity was obtainable from the extremisation of a suitable entropy functional (see also [4, 5] for a general account of this). In these approaches, arbitrary accelerating observers at a generic point of spacetime are considered. The equations of motion are obtained through the proper consideration of a key quantity, $\frac{dE}{T_H}$ (energy flux of matter crossing the horizon over horizon temperature), or variation of the Clausius entropy of the horizon, associated with the crossing of matter through local Rindler horizons. It represents the change of entropy of the “system beyond the horizon” or, in brief, of the horizon (concerning the general issue of whether patches of accelerating (or general) horizons, i.e. not only of black hole horizons, can consistently be associated with an entropy, cf. [6]).

Our aim here, is to contrast this approach with the one which obtains the equations of motion from entropy bounds. Crucial to these is the entropy dS of matter. In entropy bounds (in their generalised formulation [7] which can be thought of as subsuming many other formulations (including the first entropy bound ever formulated, the Bekenstein bound for the entropy-to-energy ratio [8]) and can be considered as a general statement of holography), the entropy dS of matter which goes through a null hypersurface is bounded by a quarter of the change of the area of the spatial cross-section of the hypersurface. The existence of such a bound is seen as remounting at the end to the existence of a lower limit to the “amount of space” to be assigned to the single bit of information, a limit of the order of the Planck length.

If one further postulates this latter feature of information is intrinsic, that is unrelated to gravity, it is then possible to *derive* the equations of motion of gravity from it. That is, from the mere existence of this fundamental discretisation of space associated with information, one can derive that matter curves spacetime, and the way it does (independent explicit implementations of this idea are [9], [10]; see also [11, 12]; it can however be seen that it was already contained in [1]; in [13, 14] and [5], a gravitational acceleration of entropic origin is also considered, but, apparently, with no intention of considering this as a *derivation* of gravity from horizon entropy). If on the contrary, on defining the limiting “amount of space” above, gravity itself intervenes, the *derivation* would turn into a *description* of gravity in terms of entropy bounds. Gravity would be one among the basic players and would no longer be reducible, at a fundamental level, to the pure thermodynamics of something else (while maintaining its emergent thermodynamic nature in the long wavelength limit).

What is behind the formulation of the entropy bounds in terms of the area/4 of suitable surfaces, is that, in case these surfaces act as (spatial cross sections of) horizons, the area/4 does coincide with horizon entropy, and this implies Einstein gravity. The approach to the equations of motion as a thermodynamic identity is independent, instead, of being the horizon entropy given by area times $\frac{1}{4}$ or by area times any other constant,

*Electronic address: pesci@bo.infn.it

or, also, given by general expressions involving, say, a dependence on curvature at each point (we are insisting, pedantically perhaps, on the expression “area times a constant” in place of “proportional to area” because the latter expression can include entropies still locally proportional to area, but not given by area times a constant, due to a dependence on the point). The results in [1] have been obtained, indeed, on assuming horizon entropy as area times a constant; but extensions of these results to more general expressions for horizon entropy have been suggested in [1] itself, and explicitly worked out in [15] for Gauss-Bonnet and general Lanczos-Lovelock models [16–18] (in a cosmological setting), in [19] (for $f(R)$ theories) and (to include any diff-invariant theory) in [20] and [4, 5], through the use of Wald’s entropy [21–23] as horizon entropy. The mentioned derivation in [2, 3] of the equations of motion from extremisation of a suitable entropy functional, lives moreover in a context far more general than Einstein gravity (Lanczos-Lovelock models).

In our bridging of the two approaches, there is thus a need to disentangle in the entropy bounds what is gravitational-dynamics dependent and what can act as a bound to matter entropy in the form of a quantity which is irrespective of the gravitational theory we are in. This is what we try to do in what follows.

The framework in which we move are arbitrary diffeomorphism invariant theories of gravity with gravitational Lagrangians depending on the metric, on the Riemann tensor, and on the derivatives of the latter of whichever order and nothing else, and with minimal coupling between matter and gravity.

II. BOUND TO THE ENTROPY OF AN ELEMENT OF MATTER AS VARIATION OF CLAUSIUS ENTROPY OF THE HOTTEST HORIZON GOING TO SWALLOW IT

In the calculation of matter fluxes through the horizon, approaches differ in the assumptions they make concerning the local Rindler horizon they are considering (for instance, Jacobson [1] and Padmanabhan [2–5]). In particular, the expansion of the null congruence which generates the horizon is assumed to be vanishing at the point under consideration in [1], while it is arbitrary in [2–5]. This difference in the assumptions, has remarkable consequences concerning the definition/interpretation of some thermodynamic potentials associated with matter fluxes through horizons, as pointed out and discussed in [24]. However it does not appear to affect the variation of Clausius entropy of the horizon, provided that a notion of Clausius entropy $\frac{dE}{T_H}$ for arbitrary bifurcate null surfaces is introduced [25].

These topics are strictly connected with present discussion. Here, however, we wish to elaborate on Clausius entropy of the horizon in its ability to bound the entropy of matter dropped through the horizon and in its independence of the gravity theory we are in, regardless of any given horizon expansion. We choose thus to consider the simplest possible case, namely the case of a patch of Rindler horizon with vanishing expansion θ . The results we shall find are applicable to any horizon locally approximable by Rindler $\theta = 0$ (as bifurcate Killing horizons are).

The Wald entropy associated with any patch of horizon, in particular to Rindler horizons with $\theta = 0$, does depend, of course, on the gravity theory we are in. The emphasis in this Section is that the variation of Clausius entropy of the horizon associated to the passage of a lump of matter through it, is instead a (gravitational) dynamics-independent quantity. The reason for this emphasis is that, given any element of matter characterized by its energy-momentum tensor T_{ab} besides its size, our aim is to construct a bound to its entropy in terms of the variation of the Clausius entropy of (a certain) horizon; a bound which thus turns out to be independent of gravitational dynamics. We do this by making use of the l^* concept, introduced earlier [26, 27].

We consider some smooth distribution of matter. We associate with it its energy-momentum tensor field T_{ab} defined in a D -dimensional (gauge [28]) spacetime ($D \geq 4$) with metric g_{ab} , the geometry of this spacetime being determined, through some field equations, by that matter and all other matter in the universe. At a generic point P in this spacetime, let k^a be a null vector, directed to the future of P . We then consider a local inertial frame around P , with coordinates X^a , such that P is at $X^a = 0$, and chosen in such a way as to have k^a given by (1,1) in the plane (T, X) . The null curve (X, X) has affine parameter X and tangent k^a . We consider a small piece of matter around P . The element of matter has a generic velocity in the local inertial frame. Its flow across any patch of surface is determined by the velocity vector, which has an intrinsic meaning independent of the gravity theory which determined a given spacetime as solution (and back-reaction effects, due to shrinking of the area while the element of matter goes through, are second order effects with respect to this in the flow).

Let us next consider the local Rindler frame [1, 4] associated with an observer accelerating along X with acceleration κ , which is at rest with respect to the local inertial frame at Rindler time $t = 0$. We know that the metric can be written as $ds^2 = -(kx)^2 dt^2 + dx^2 + dx_\perp^2 \equiv -N^2 dt^2 + dx^2 + dx_\perp^2$ with obvious notation, with x^a Rindler coordinates and dx_\perp^2 the Euclidean metric in the $(D - 2)$ -plane. In these coordinates, the accelerating observer \mathcal{A} is, at $t = 0$, at $x = \frac{1}{\kappa}$.

The Killing vector $\xi_{(t)}^a \equiv \left(\frac{\partial}{\partial t}\right)^a$, corresponding to time translation invariance, null on the horizon and orthogonal to it, is what is needed to express the mass M of the horizon (of the “system beyond the horizon”),

attributed to it by the accelerating observer. The condition $\xi_{(t)}^a \xi_{(t)a} = -N^2 = 0$, that is $x = 0$, or $T \pm X = 0$, gives the location of the horizon. To this, the accelerating observer, which sees herself in a thermal bath, assigns a temperature $T_H = \frac{\kappa}{2\pi}$. For the entire construction to be consistent, the size $L_R \approx \frac{1}{\kappa}$ associated to the Rindler description must be much smaller than the size L_I on which the local inertial frame approximation works, that is $L_R \ll L_I \approx \frac{1}{\sqrt{\mathcal{R}}}$, where \mathcal{R} is the magnitude of a typical component of the curvature tensor. We assume to have chosen κ large enough such that this is the case.

Let us first review how it goes in the standard approach, that the variation of the Clausius entropy of the horizon associated with the passage of an element of matter is independent of the gravitational dynamics. For this, we have to obtain the energy flux dE entering in the expression $\frac{dE}{T_H}$. It coincides (since the horizon is orthogonal to $\xi_{(t)}^a$) with the increment in mass dM of the horizon. Standard calculation gives

$$dE = T_{ab} \xi_{(t)}^a k^b A dX, \quad (1)$$

where A is the area of the cross-section of the element of matter in the $(D-2)$ -plane, and dX its size in X -coordinate. This quantity depends on the characteristics of the lump of matter (T_{ab} , A , dX) and on the geometric characteristics of the horizon ($\xi_{(t)}^a$, k^b); in no way can it depend on the equations of motion of gravity, whatever they are, having g_{ab} as solution associated with the assigned distribution of matter. The same can be said of the associated Clausius entropy variation of the horizon, since

$$\frac{dE}{T_H} = \frac{1}{T_H} T_{ab} \xi_{(t)}^a k^b A dX, \quad (2)$$

and T_H is given with the field $\xi_{(t)}^a$. Thus, the expression of the amount of Clausius entropy of the horizon associated with the passage of a lump of matter through it, displays in an obvious way the independence of this quantity with respect to gravitational dynamics.

Let us now stay with the element of matter, and consider any possible local Rindler horizon with any direction and any temperature, i.e. the horizon perceived by any possible accelerating observer sitting instantaneously where matter is, and about to absorb it. We ask about the relation between the entropy dS of the lump of matter and the variation of the Clausius entropy of any of these horizons. This corresponds to considering the situation depicted in Fig. 1, with $\{X^a\}$ now being the local frame of matter and the element of matter located at $X = \frac{1}{\kappa}$ at $t = 0$, and asking about the relation between dS and $\frac{dE}{T_H}$.

We evaluate dS from Gibbs-Duhem relation (for a 1-component thermodynamic system), in its local form

$$\rho = s\theta - P + \mu n, \quad (3)$$

a relation which merely expresses the first law of thermodynamics joined to the request of (local) extensivity of pertinent thermodynamic potentials (cf. [29]). Here, ρ , s , θ , P , μ and n are respectively local energy density, entropy, temperature, pressure, chemical potential and number density of the element of matter. As we are concerned with gravity, the rest-mass energy is thought here to be included in ρ ; we know this means that this same energy must be thought of as also included in μ (cf., for example, [30] (p. 155)). We have

$$dS = s dV_{prop} = \frac{\rho + P - \mu n}{\theta} dV_{prop}, \quad (4)$$

where dV_{prop} is the proper volume of the element of matter.

In the calculation of $\frac{dE}{T_H}$, we explicitly choose $T_{ab} = (\rho + P)u_a u_b + P g_{ab}$ (u^a is the velocity vector of the element of matter), that is we choose the stress-energy tensor of an ideal fluid, in the assumption that this case is general enough to show what is here happening concerning the relation between dS and $\frac{dE}{T_H}$, without, unnecessary at this stage, further complications. In (2), all we need to know is $\xi_{(t)}^a$ on any given horizon. From $\xi_{(t)}^a \rightarrow \kappa X k^a$ on the horizon (cf., for instance, [4]), we get

$$\frac{dE}{T_H} = \frac{1}{T_H} T_{ab} k^a k^b dV_{prop}, \quad (5)$$

and thus

$$\frac{dE}{T_H} = \frac{\rho + P}{T_H} dV_{prop}. \quad (6)$$




FIG. 1: (Piece of) worldline of an element of volume at $X = \frac{1}{\kappa}$, at rest in the inertial frame (see text).

Comparing dS with $\frac{dE}{T_H}$, we see amounts to compare $s = \frac{\rho+P-\mu n}{\theta}$ with $\frac{\rho+P}{T_H}$. In this regard, we notice that what we are talking about is an element of matter of some proper size l , with temperature and other thermodynamic potentials assigned, which, initially, i.e. at $t = 0$ and at a proper distance $D = \frac{1}{\kappa}$ from the horizon, is assumed to be entirely ahead of the horizon. This means $l \leq 2D = \frac{2}{\kappa}$.

The circumstances can be conveniently described in terms of the length l^* mentioned above, characteristic of the given local thermodynamic conditions. The meaning of l^* is that, for whatever thermodynamic system with given (local) thermodynamic conditions, its size Δ cannot be smaller than the l^* for these conditions. More precisely,

$$\Delta \geq l^* \equiv \frac{1}{\pi} \frac{s}{\rho + P} = \frac{1}{\pi\theta} \left(1 - \frac{\mu n}{\rho + P} \right), \quad (7)$$

where the last equality follows from Gibbs-Duhem relation. In practice this means that, if we insist in saying that a system has certain given values of ρ , s , and P , we have to realize that that system cannot have size smaller than l^* as given in (7).

Inequality (7) is not universally proven, it is a conjecture. It is however likely to be obeyed by any matter system, as an expression of the unavoidable quantum indeterminacy of the constituent particles, i.e. its reason rests on that it seems to be simply a manifestation, or reformulation, of the uncertainty principle. As such, it can be considered as a universal feature of any thermodynamic system, and it is moreover a basic expression of quantum mechanics alone (i.e. without reference to gravity, meaning with reference to Minkowski limit only). In [26, 27] an analysis is made on how the inequality (7) arises in some relevant and calculable systems.

Applying this here, we get

$$s \leq \pi l(\rho + P) \leq \frac{2\pi}{\kappa}(\rho + P) = \frac{\rho + P}{T_H}, \quad (8)$$

and so

$$dS \leq \frac{dE}{T_H}, \quad (9)$$

where the equality is reached when $l = l^*$ and $l = \frac{2}{\kappa}$.

The possibility of fulfilling the condition $l = l^*$ is tied to the nature of the matter under consideration. For certain material media the equality can indeed in principle be reached; for example this happens for ultrarelativistic systems with $\mu = 0$, e.g. a blackbody photon gas, when the size l of the system is made very small,

up to the limit of compatibility, in view of quantum uncertainty, with the assigned values of thermodynamic potentials [27]. In general media, even when this quantum limit is reached, l is still much larger than l^* . For generic choices of the size of the element of matter, we will have $l \gg l^*$ even for photon gases.

The condition $l = \frac{2}{\kappa}$ deals with the location of the horizon which absorbs the element of matter. When $l = \frac{2}{\kappa}$ the horizon just starts to absorb, that is the element of matter is just ahead of the horizon, at the limit of the beginning of the absorption; and it is $l < \frac{2}{\kappa}$ otherwise, i.e. when there is still some path before absorption begins. The eventuality $l > \frac{2}{\kappa}$ is not allowed, since it is incompatible with the assumption that the system is entirely ahead of the horizon at start; a system with $l > \frac{2}{\kappa} = \frac{1}{\pi T_H}$, where T_H is the temperature of the horizon about to absorb it, has necessarily a part of itself beyond the reach of the Rindler coordinates of the accelerating observer located where matter is and instantaneously at rest with respect to it, whom perceives that horizon, so that this part of the system results already absorbed by the horizon.

We have thus shown the following.

Proposition. Given an element of matter of size l and energy-momentum tensor T_{ab} , its entropy dS is bounded from above by the quantity $\frac{dE}{T_H}$, given by (5), of a Rindler horizon at temperature $T_H = \frac{1}{\pi l}$ as perceived by an accelerating observer sitting instantaneously where matter is, with k^a the tangent vector to the generators of the horizon, i.e. by the variation of Clausius entropy of a Rindler horizon at the perceived temperature $T_H = \frac{1}{\pi l}$, which engulfs the element of matter.

The limiting value can in principle be reached: this happens for ultrarelativistic systems with $\mu = 0$, of very small size, pushed to the (quantum) limit of compatibility with the assigned thermodynamic conditions for the element of matter. The bound can thus also be considered as the supremum among the dS for all possible material media, for given dE and l . Thus considered, it is manifest that the bound is a quantity inherent to matter, and is irrespective of the gravity theory, in agreement with what we already know since, as seen, the quantity $\frac{dE}{T_H}$ is independent of the gravity theory.

III. THE GENERALIZED COVARIANT ENTROPY BOUND EXTENDED TO GENERAL THEORIES OF GRAVITY

In the descriptions-derivations of the equations of motion as a thermodynamic identity, the equality

$$\frac{dE}{T_H} = dS_W \quad (10)$$

between the Clausius entropy variation of the horizon to the left, and the Wald entropy variation of horizon to the right (coinciding with the Bekenstein-Hawking entropy variation dS_{BH} in Einstein theory) associated with the crossing of the horizon by some element of matter is stressed as entailing the equations of motion for the gravitational field, i.e. how matter curves geometry. Indeed, Wald's entropy S_W is a prescription which, on using the equations of motion, gives the amount of entropy associated with a patch of horizon of some $(D-2)$ -dimensional cross-sectional area A , the amount of entropy being different for different gravity theories. The l.h.s. is clearly independent of the gravity theory, as can be seen, for example, on noting that any bifurcate horizon can be approximated at a point by a Rindler horizon, and applying the discussion of previous Section. The r.h.s. must also be independent of the gravity theory. Equality (10) then gives how much the area A must shrink, due to some passage of matter, this depending on the expression of S_W . For each expression of S_W this gives the equations of motion for the theory which has that expression as its Wald's entropy of the horizon.

Using this, the bound discussed in the previous Section becomes, in each specific theory of gravity, a bound in terms of the Wald entropy of that theory. From (9) and (10), in fact, we get

$$dS \leq dS_W. \quad (11)$$

According to previous Section we are considering this relation in terms of a patch of Rindler horizon at P with tangent k^a to the generators and temperature $T_H = \frac{\kappa}{2\pi}$. We can however think of dS_W as referring to a generic bifurcate Killing horizon H at P with normal ξ_H , tangent k_H^a to the generators and surface gravity κ_H , on calculating it through the approximating Rindler patch at P with $\xi_{(t)} = \xi_H$, $k^a = k_H^a$ and $\kappa = \kappa_H$.

In D -dimensional Einstein gravity S_W coincides with S_{BH} , and (11) becomes

$$dS \leq dS_{BH} = \frac{dA}{4}. \quad (12)$$

But, this is the generalized covariant entropy bound [7] (GCEB), as applied to the patch of $(D-2)$ -hypersurface of area A coinciding with the $(D-2)$ -section of a patch of a horizon and having k_H^a as the null field orthogonal

to it. Inequality (11) thus generalizes the GCEB to a form which applies in any metric theory of gravity, with the generalization being in that $\frac{dA}{4}$, with A the $(D-2)$ -area of a patch B of $(D-2)$ -hypersurface with an orthogonal null field k^a , is replaced by dS_W , thought to be the Wald entropy of the patch of the sheet of null hypersurface generated by null geodesics which start at B with tangent k^a .

IV. l^* FOR HORIZONS

The scale l^* has been considered in the past for conventional matter only [26, 27]. The discussion of Section II gives the opportunity of defining a value l_H^* for it for horizons. The definition of l^* given in (7) can evidently be put also in the form

$$l^* \equiv \frac{1}{\pi} \frac{s dV_{prop}}{(\rho + P)dV_{prop}} = \frac{1}{\pi} \frac{dS}{dE}. \quad (13)$$

This is the l^* of the matter which is going to be swallowed by a horizon. The matter has an entropy content dS in some proper volume $dV_{prop} = Al$ and gives an energy contribution dE to the horizon, which has a given temperature T_H . Now, if we use matter for which the limit $l = l^*$ can be reached, and we choose $l = l^*$, and if we furthermore assume that our choice of the thermodynamic parameters of matter is such that $l = l^*$ is the maximum value allowed for the matter to be absorbed in one bite by the given horizon, i.e. $l = \frac{1}{\pi T_H}$, we know that in (9) the equality holds, and we are thus allowed to write $\frac{dE}{T_H}$ for dS in (13). But, the member to the right in (13) is now given in terms of quantities which refer to the horizon alone. This suggests

$$l_H^* \equiv \frac{1}{\pi} \frac{dE/T_H}{dE} = \frac{1}{\pi T_H} = \frac{1}{\pi} \left(\frac{s}{\rho + P} \right)_{l=l^*=\frac{1}{\pi T_H}}, \quad (14)$$

where the subscript reminds us that the term in the round brackets must be evaluated for material media for which the choice $l = l^* = \frac{1}{\pi T_H}$ is allowed. We said above that a blackbody photon gas, for which $l^* = \frac{1}{\pi\theta}$ (cf. (7)), does the job in that it allows for $l = l^*$ and we get $l = \frac{1}{\pi T_H}$ provided the temperature of the gas equals the horizon temperature, $\theta = T_H$. l_H^* is thus the order of the wavelength of blackbody photons at the temperature T_H of the thermal bath perceived by the accelerating observer. This quantity is supposed to conveniently characterise the behaviour of a horizon from a pure quantum mechanical point of view, meaning in the Minkowski limit (and we know horizons do not imply we leave this limit). As such, it does not contain any information on gravity, and is well defined also in a context for which, spacetime is purely Minkowskian (cf. [27]). It is therefore a concept clearly distinct from the Planck scale l_P (and, in general, is enormously larger than the latter).

For a generic bifurcate Killing horizon, its l_H^* can be defined in an obvious way through the Rindler horizon approximating it at a point. For systems collapsing to form black holes in Einstein gravity with temperature T_H in asymptotically-flat spacetimes, for example, the photon gas (or ultrarelativistic matter) above, which gives $l = l^* = \frac{1}{\pi T_H}$, is a blackbody and must have temperature $\theta = T_H$, as measured by distant observers (as can be verified from locally Rindler approximating the horizon); i.e., it turns out it is just their Hawking radiation. For the Schwarzschild black hole we get

$$l_H^* = \frac{1}{\pi T_H} = \frac{8\pi M}{\pi} = 8M = 4R, \quad (15)$$

where M and R are black hole mass and radius.

V. CONCLUDING REMARKS

We have seen the limiting thermodynamical scale l^* of matter can be used to show that entropy of any element of matter is bounded by the variation of Clausius entropy of a suitable Rindler horizon absorbing it. From this, a form of the GCEB, valid for general metric theories of gravity, has been introduced, and the l^* for horizons has been defined.

Using the perspective according to which the entropy of any patch of horizon is always (i.e. for any diff-invariant theory of gravity) given by a quarter of area in units of effective coupling [31], from (11) we get

$$dS \leq \frac{dA}{4G_{eff}}, \quad (16)$$

where the effective coupling G_{eff} , defined as in [31], is explicitly reported, and, again, A is the area of a patch of the boundary surface. G_{eff} depends in general on the point in spacetime, and G_{eff} equals Newton's constant, $G_{eff} = \text{constant} = G_N$, for Einstein gravity. Formula (16) is a further (local) expression of the GCEB generalized to general theories of gravity. When we change the gravity theory, the quantity $\frac{dE}{T_H}$ and thus $\frac{dA}{4G_{eff}}$ remains unaffected. What changes, are dA and G_{eff} individually. A gravity theory with a gravitational coupling, say, larger than that of Einstein's theory ($G_{eff} > G_N$), will give a stronger focusing (a larger dA), stronger by the amount exactly needed to leave $\frac{dA}{4G_{eff}}$ invariant.

The fact that the black hole horizon has the same l^* as its Hawking radiation, can be interpreted as suggesting that in some respects it behaves like the Hawking radiation itself. This radiation is, after all, all that a distant observer can see of the black hole. This thermodynamic fact, could provide some further understanding of the dynamic behaviour of black holes. The dynamic relaxation times of perturbed black holes (for a review see [32]), could be seen as an example of this kind. The argument for this, goes as follows. A universal lower limit $\tau_0(\theta) \equiv \frac{1}{\pi\theta}$ to the relaxation times τ of systems at temperature θ , has been introduced in [33] from thermodynamics and quantum information theory. As far as black holes can be considered thermodynamic objects, we would expect that $\tau_0(T_H)$ could also limit the dynamic relaxation times of perturbed black holes with horizon temperature T_H . Could this be inspected through the use of l^* for horizons? Systems consisting of blackbody radiation at temperature θ actually obey this limit, and, shaped as thin layers, they exhibit, when sufficiently thin, relaxation times τ approaching it, $\tau = \tau_0(\theta)$ (in the limit $l \rightarrow l^*(\theta)$, l being their thickness, limit that, as mentioned above, for blackbody radiation can be reached) [34]. Now, the imprint on any radiation, of the variation of the properties of the black hole cannot have any evolution more rapid than that given by this limit with $\theta = T_H$. That means, the distant observer cannot see variations of the black hole on a characteristic scale smaller than this, i.e. the relaxation times of black holes cannot be smaller than this limit: $\tau \geq \tau_0 = \frac{1}{\pi T_H}$. This, namely that black holes comply with the bound, is precisely what is obtained by analytical and numerical evaluations [33]. In [33] it is found moreover that, when black holes are nearly extremal, i.e. in the $T_H \rightarrow 0$ limit, they saturate the bound. According to the perspective above, near-extremal black holes could thus be regarded as equivalent to a thickness $l = l_H^* = \frac{1}{\pi T_H}$ of their own Hawking radiation, from a far observer point of view.

A last comment concerns the particular case of matter for which $P = -\rho$, i.e. matter mimicking the effects of a cosmological constant term in the equations of motion. For this matter, the Gibbs-Duhem relation (3) becomes

$$s\theta + \mu n = 0, \quad (17)$$

where s , with its informational meaning, and θ and n should be regarded as intrinsically non-negative. We have two possibilities:

- i) $s > 0$; implying $\mu < 0$;
- ii) $s = 0$; implying $\mu n = 0$.

Looking at (6), both cases give $\frac{dE}{T_H} = 0$, and we do not get an increment of Clausius entropy of the horizon when such matter crosses it. In case (i), this gives a violation of (9), and thus of (11). In l^* terminology, the situation is characterized by $l^* = \infty$ in (7), and thus we see that we can never have $l \geq l^*$ for this matter for any l ; that is, matter of this kind is incompatible with the assumption for it to be initially completely ahead of the horizon. The basic assumption itself that matter can be described in terms of local quantities (ρ, s, \dots) in a volume dV_{prop} , a pre-requisite for writing the local form of the Gibbs-Duhem relation (3), is put into question; the condition itself $s > 0$ becomes of doubtful meaning in that s is found to be ill-defined. In view of this, we cannot claim that any matter of this kind would produce a violation of entropy bound (11); what we could claim is that the bound could hold true even in presence of this “matter”, since the real point is that surely our description of it in local terms is no longer adequate (this perhaps offers a different perspective on the issue of the effect of any cosmological constant term on entropy bounds as tackled in [35]). We could speculate that, if we believe in quantum mechanics as providing l^* , the case $s > 0$ would correspond to “cosmological matter” which would intrinsically be completely delocalised ($l^* = \infty$). If we think of constituents for it, they should be completely delocalised, and maximally entangled.

Case (ii) gives $l^* = \frac{0}{0}$, that is l^* is undetermined without further input. Any finite value of l^* would allow inequality (9) to be satisfied irrespective of l , i.e. even when $l < l^*$. This would suggest, for overall consistency, $l^* = 0$ for case (ii). It would correspond to matter completely localised, with no entanglement, with infinite energy and pressure (from quantum indeterminacy), and no entropy.

Acknowledgments. I thank Giovanni Venturi for the careful reading of the manuscript.

-
- [1] T. Jacobson, “Thermodynamics of spacetime: the Einstein equation of state”, *Phys. Rev. Lett.* **75** (1995) 1260, gr-qc/9504004.
 - [2] T. Padmanabhan, A. Paranjape, “Entropy of null surfaces and dynamics of spacetime”, *Phys. Rev. D* **75** (2007) 064004, gr-qc/0701003.
 - [3] T. Padmanabhan, “Dark energy and gravity”, *Gen. Rel. Grav.* **40** (2008) 529, arXiv:0705.2533.
 - [4] T. Padmanabhan, *Gravitation: Foundations and frontiers* (Cambridge Univ. Pr., Cambridge UK, 2010).
 - [5] T. Padmanabhan, “Thermodynamical aspects of gravity: new insights”, *Rept. Prog. Phys.* **73** (2010) 046901, arXiv:0911.5004.
 - [6] T. Jacobson, R. Parentani, “Horizon entropy”, *Found. Phys.* **33** (2003) 323, gr-qc/0302099.
 - [7] É.É. Flanagan, D. Marolf and R.M. Wald, “Proof of classical versions of the Bousso entropy bound and of the generalized second law”, *Phys. Rev. D* **62** (2000) 084035, hep-th/9908070.
 - [8] J.D. Bekenstein, “Universal upper bound on the entropy-to-energy ratio for bounded systems”, *Phys. Rev. D* **23** (1981) 287.
 - [9] T. Jacobson, “Gravitation and vacuum entanglement entropy”, arXiv:1204.6349 (2012).
 - [10] E.P. Verlinde, “On the origin of gravity and the laws of Newton”, *JHEP* **1104** (2011) 029, arXiv:1001.0785.
 - [11] A. Pesci, “Gravity from the entropy of light”, *Class. Quantum Grav.* **28** (2011) 045001, arXiv:1002.1257.
 - [12] A. Pesci “The existence of a minimum wavelength for photons”, arXiv:1108.5066 (2011).
 - [13] T. Padmanabhan, “Entropy of static spacetimes and microscopic density of states”, *Class. Quantum Grav.* **21** (2004) 4485, gr-qc/0308070.
 - [14] T. Padmanabhan, “Equipartition of energy in the horizon degrees of freedom and the emergence of gravity”, *Mod. Phys. Lett. A* **25** (2010) 1129, arXiv:0912.3165.
 - [15] R.-G. Cai, S.P. Kim, “First law of thermodynamics and Friedmann equations of Friedmann-Robertson-Walker universe”, *JHEP* **0502** (2005) 050, hep-th/0501055.
 - [16] C. Lanczos, “Electricity as a natural property of Riemannian geometry”, *Rev. Mod. Phys.* **39** (1932) 716.
 - [17] C. Lanczos, “A remarkable property of the Riemann-Christoffel tensor in four dimensions”, *Annals Math.* **39** (1938) 842.
 - [18] D. Lovelock, “The Einstein tensor and its generalizations”, *J. Math. Phys.* **12** (1971) 498.
 - [19] E. Elizalde and P.J. Silva, “F(R) gravity equation of state”, *Phys. Rev. D* **78** (2008) 061501, arXiv:0804.3721.
 - [20] R. Brustein, M. Hadad, “The Einstein equations for generalized theories of gravity and the thermodynamic relation $\delta Q = T\delta S$ are equivalent”, *Phys. Rev. Lett.* **103** (2009) 101301, Erratum-ibid. **105** (2010) 239902, arXiv:0903.0823.
 - [21] R.M. Wald, “Black hole entropy is Noether charge”, *Phys. Rev. D* **48** (1993) 3427, gr-qc/9307038.
 - [22] T. Jacobson, G. Kang, R.C. Myers, “On black hole entropy”, *Phys. Rev. D* **49** (1994) 6587, gr-qc/9312023.
 - [23] V. Iyer and R.M. Wald, “A comparison of Noether charge and Euclidean methods for computing the entropy of stationary black holes”, *Phys. Rev. D* **52** (1995) 4430, gr-qc/9503052.
 - [24] D. Kothawala, “The thermodynamic structure of Einstein tensor”, *Phys. Rev. D* **83** (2011) 024026, arXiv:1010.2207.
 - [25] V. Baccetti, M. Visser, “Clausius entropy for arbitrary bifurcate null surfaces”, arXiv:1303.3185.
 - [26] A. Pesci, “From Unruh temperature to the generalized Bousso bound”, *Class. Quantum Grav.* **24** (2007) 6219, arXiv:0708.3729.
 - [27] A. Pesci, “On the statistical-mechanical meaning of the Bousso bound”, *Class. Quantum Grav.* **25** (2008) 125005, arXiv:0803.2642.
 - [28] C. Rovelli, *Quantum gravity* (Cambridge Univ. Pr., Cambridge UK, 2004).
 - [29] J. Oppenheim, “Thermodynamics with long-range interactions: from Ising models to black-holes”, *Phys. Rev. E* **68** (2003) 016108, gr-qc/0212066.
 - [30] K. Huang, *Statistical Mechanics* (J. Wiley & Sons, New York, 1987).
 - [31] R. Brustein, D. Gorboson, M. Hadad, “Wald’s entropy is equal to a quarter of the horizon area in units of the effective gravitational coupling”, *Phys. Rev. D* **79** (2009) 044025, arXiv:0712.3206.
 - [32] H.-P. Nollert, “Quasinormal modes: the characteristic ‘sound’ of black holes and neutron stars”, *Class. Quantum Grav.* **16** (1999) R159.
 - [33] S. Hod, “Universal bound on dynamical relaxation times and black-hole quasinormal ringing”, *Phys. Rev. D* **75** (2007) 064013, gr-qc/0611004.
 - [34] A. Pesci, “A note on the connection between the universal relaxation bound and the covariant entropy bound”, *Int. J. Mod. Phys. D* **18** (2009) 831, arXiv:0807.0300.
 - [35] J.-W. Lee, “Zero Cosmological Constant and Nonzero Dark Energy from Holographic Principle”, *J. Korean Phys. Soc.* **63** (2013) 1088, arXiv:1003.1878.